arXiv:1209.6418v1 [math.AG] 28 Sep 2012

Colliot-Thélène's conjecture and finiteness of u-invariants

Max Lieblich R. Parimala V. Suresh

October 1, 2012

Introduction

Let X an excellent Noetherian regular scheme of dimension 2 which is projective over an affine Dedekind scheme Spec(A) and K the function field of X. Let l be a prime which is a unit on X (i.e. unequal to characteristic of the residue field of any point in X). The work of Saltman on the division algebras over surfaces ([S1], [S3]) imply that given an element α in the l-torsion of the Brauer group of K, there exist elements $f, g, \in K^*$ such that $\alpha \otimes K(\sqrt[l]{f}, \sqrt[l]{g})$ is unramified on a regular proper model of $K(\sqrt[l]{f}, \sqrt[l]{g})$ over A (cf. [B]). If K is the function field of a curve over a p-adic field and l a prime not equal to p, the unramified Brauer group on a regular proper model over $\mathbf{Z}_p$ is zero. Hence, for $\alpha \in {}_lBr(K)$, index of α divides l^2.

In this paper, we split the ramification of division algebras on surfaces in a more general setting without any assumption on the characteristic of the residue fields of points of the scheme. More precisely, we prove the following (cf. Theorem 2.9):

Theorem. Let X be an excellent regular integral scheme of dimension 2 and K its function field. Suppose that $\mathrm{char}(K) = 0$ and for every codimension one point x of X, if the characteristic of the residue field $\kappa(x)$ at x is p, then, $[\kappa(x) : \kappa(x)^p] = p$ (i.e. p-dimension of $\kappa(x)$ is 1). If α is an element in the p-torsion of the Brauer group of K, then there exists $f, g, h \in K^*$ such that $\alpha \otimes K(\sqrt[p]{f}, \sqrt[p]{g}, \sqrt[p]{h})$ is unramified on a regular model Y of K proper over X.

The above theorem leads to the following results on splitting ramification of exponent p division algebras over function fields of curves over p-adic fields and number fields (cf. Corollaries 2.10 and 2.11).

Corollary. If k is a p-adic field and K a function field of a curve over k, then for every element $\alpha \in {}_pBr(K)$, index(α) divides period$(\alpha)^3$.

Corollary. Let k be a number field, S the ring of integers in k, and K the function field of a curve over k. Let $\alpha \in Br(K)$ be a p-torsion element.There is an explicit degree p^3 extension L of K such that $\alpha \otimes L$ is the restriction of a class in the Brauer group of a regular proper model of L over S.

Thus the study of the period-index problem for such function fields is reduced to a corresponding problem for unramified Brauer classes. Recall the following conjecture of Colliot-Thélène:

CT-Conjecture([CT1]): If Y is a smooth projective geometrically connected variety over a global field k then the Brauer-Manin obstruction to the existence of 0-cycles of degree 1 is the only obstruction.

Using a moving lemma for 0-cycles, we extend results of [L2] to show the following (cf. Theorem 3.3).

Theorem. Let k be a number field and K the function field of a curve over k. Let X be a regular proper model of K over the ring of integers in k. If the CT-Conjecture holds then for any $\alpha \in \mathrm{Br}(X)$, the ind$(\alpha)$ divides per$(\alpha)^2$.

Combining this with the results described above on splitting ramification, we deduce the following corollary (cf. Theorem 4.1).

Corollary. Let K be a function field in one variable over a number field. The CT-conjecture implies that for any $\alpha \in Br(K)$, ind(α) divides period$(\alpha)^5$.

This together with a result on degree 3 Galois cohomology groups ([Su]) leads to the final result of the paper (cf. Theorem 4.4).

Theorem. If the CT-Conjecture holds then the function field of a curve over a totally imaginary number field has finite u-invariant.

Acknowledgements: We thank D. Saltman for generously sharing with us his notes on the Brauer group of complete discrete valuated fields in in equicharacteristic. The first author is partially supported by the Sloan Foundation and National Science Foundation CAREER grant DMS-1056129. The second author is partially supported by National Science Foundation grant DMS-1001872. All authors are grateful to the American Institute of Mathematics, where this project was started as part of the workshop "Deformation theory, patching, quadratic forms, and the Brauer group" in January of 2012.

1. Brauer group of discretely valued fields

Let (K, ν) be a complete discrete valued field with ring of integers R, maximal ideal m and residue field κ. Suppose that $\mathrm{char}(K) = 0$, $\mathrm{char}(\kappa) = p > 0$ and that K contains a primitive p^{th} root of unity ζ. Write $N = \nu(p)p/(p-1)$. Since

$$(p-1)\nu(\zeta - 1) = \nu(p),$$

N is a positive integer divisible by p. Finally, for $i \geq 0$, let

$$U_i = \{u \in R^* \mid x \equiv 1 \bmod m^i\}.$$

The following assumption about the residue field will play a key role throughout this paper.

Assumption ($*$): $[\kappa : \kappa^p] = p$.

Given $a, b \in K^*$, let $(a, b) \in {}_p\mathrm{Br}(K)$ be the class of the cyclic K-algebra defined by the relations

$$x^p = a$$
$$y^p = b$$
$$xy = \zeta yx.$$

Let $br(K)_0 = {}_p\mathrm{Br}(K)$ and for $i \geq 1$, let

$$br(K)_i \subset br(K)_0$$

be the subgroup generated by cyclic algebras (u, a) with $u \in U_i$ and $a \in K^*$. Since K is complete, $br(K)_n = 0$ for $n > N$ ([CT2], 4.1.3 and [MS]). In this section we recall a few basic facts about ${}_p\mathrm{Br}(K)$ and $br(K)_n$ due to Kato and Saltman.

Lemma 1.1. (Kato [K], Thm. 2) Let

$$\alpha \in br(K)_m \setminus br(K)_{m+1}$$

with $0 \leq m \leq N$. Let $\pi \in R$ be a parameter and $f \in R^*$ with $\overline{f} \notin \kappa^p$.

(a) If $m = 0$, then

$$\alpha \equiv (u, \pi) \pmod{br(K)_1}$$

for some $u \in R^*$ with $\overline{u} \notin \kappa^p$.

(b) If m is coprime to p, then there exists $x \in R^*$ such that

$$\alpha \equiv (1 + \pi^m x, f) \pmod{br(K)_{m+1}}.$$

(c) If $0 < m < N$ and m a multiple of p, then

$$\alpha \equiv (1 + \pi^m x, \pi) \pmod{br(K)_{m+1}}$$

for some $x \in R^*$ with $\overline{x} \notin \kappa^p$.

(d) Let $b = \zeta - 1$. If $m = N$ then

$$\alpha = (1 + xb^p, f) + (1 + x'b^p, \pi)$$

for some $x, x' \in R$. Further $(1 + xb^p, f)$ is unramified at ν.

Proof. Let $K_2(\kappa)$ be the Milnor K-group and

$$k_2(\kappa) = K_2(\kappa)/pK_2(\kappa).$$

The group $k_2(\kappa)$ is isomorphic to a subgroup of Ω^2_κ (cf. [CT], 3.0). Since (by Assumption $(*)$) we have

$$[\kappa : \kappa^p] = p,$$

we know that

$$\Omega^2_\kappa = 0$$

and hence $k_2(\kappa) = 0$. Define a map

$$\kappa^*/\kappa^{*p} \to br(K)_0/br(K)_1$$

by

$$\lambda \mapsto (\tilde{\lambda}, \pi),$$

where for any $\lambda \in \kappa$, $\tilde{\lambda} \in R$ is a lift. By ([K], Thm. 2, cf. [CT], 4.3.1), this map is an isomorphism. Hence every element in $br(K)_0 \setminus br(K)_1$ is equivalent to (u, π) modulo $br(K)_1$ for some $u \in R^*$ with $\overline{u} \notin \kappa^p$, establishing part (a).

Let us prove part (b). Suppose $0 < m < N$ is coprime to p. Define a map

$$\Omega_\kappa \to br(K)_m / br(K)_{m+1}$$

by

$$x\frac{dy}{y} \mapsto (1 + \pi^m \tilde{x}, \tilde{y}),$$

where

$$x, y \in \kappa^*$$

are arbitrary elements with lifts

$$\tilde{x}, \tilde{y} \in R^*.$$

By ([K], Thm. 2), this map is an isomorphism. Since $\overline{f} \notin \kappa^p$, we have $\kappa = \kappa^p(\overline{f})$. In particular, since the (absolute) differential vanishes on κ^p, every element in Ω^1_κ has the form

$$\omega = \sum_{i=1}^{p-1} \alpha_i \overline{f}^i \frac{d\overline{f}}{\overline{f}} \tag{1}$$

for $\alpha_i \in \kappa^p$. Thus every element of $br(K)_m / br(K)_{m+1}$ is the image of an element of the form $(1 + x\pi^m, f)$ for some $x \in R^*$.

Now we will prove part (c). Let $1 < m < N$ be divisible by p. Define a map

$$\kappa / \kappa^p \to br(K)_m / br(K)_{m+1}$$

by

$$\lambda + \kappa^p \to (1 + \pi^m \tilde{\lambda}, \pi),$$

where for $\lambda \in \kappa$ we have chosen a lift $\tilde{\lambda} \in R$. Using (1) above and the fact that d is κ^p-linear, we see that every element of Ω^1_κ is closed. Thus, the first summand in ([K], map (iii) immediately preceding Thm. 2) is trivial. The map described above is just the composition with the remaining summand of ([K], map (iii) before Thm. 2) with the natural map $u_2^{(n)} / u_2^{(n+1)} \to br_n(K)$

(in the notation of [K]). Hence every non-zero element in $br(K)_m/br(K)_{m+1}$ is the image of an element of the form $(1+\pi^m x, \pi)$ for some $x \in R^*$ with $\overline{x} \notin \kappa^p$.

To prove part (d) (the case $n = N$), let $b = \zeta - 1$. By ([K], Thm. 2) or ([CT2], Thm. 4.3.1(d)) and using (1) above, the element α can be written in the desired form

$$\alpha = (1 + xb^p, f) + (1 + x'b^p, \pi).$$

It remains to show that the first summand is unramified (i.e., split by an unramified extension of K). If $x \in \pi R$ then the first summand lies in $br_{N+1}(K) = 0$ and is unramified.

Thus, we may assume that $x \in R^*$, so that $\nu(x) = 0$. We will show that in this case the field

$$L = K(\sqrt[p]{1 + xb^p})$$

is unramified over K; since this splits the algebra $(1 + xb^p, f)$, this will complete the proof.

Let

$$\theta = \sqrt[p]{1 + xb^p}$$

and

$$d = \theta - 1.$$

Let $\tilde{\nu}$ be the valuation on L extending ν on K. Since $1 + xb^p$ is not a pth power, the minimal polynomial of θ is

$$g(z) = (z+1)^p - 1 - xb^p = 0.$$

In particular, the norm of θ is $-xb^p$, so

$$\tilde{\nu}(\theta) = \frac{1}{p}\nu(-xb^p) = \nu(b)$$

by our assumption that $x \in R^*$. This means that $d = wb$ for some unit w in the ring of integers in L. It is easy to see that $\overline{w}^p - \overline{w} = \overline{x}$. Hence the L/K is an unramified extension, as desired. □

Corollary 1.2. Every element $\alpha \in {}_p\mathrm{Br}(K)$ is of the form $(g, f) \cdot (h, \pi)$ for some $g, h \in R^*$.

Proof. Let $0 \leq n \leq N$. Then, by (1.1), every element in $br(K)_n$ is equivalent to $(g_n, f)(h_n, \pi)$ modulo $br(K)_{n+1}$ for some $g_n, h_n \in R^*$. Since $br(K)_n = 0$ for $n > N$, the corollary follows. □

Lemma 1.3. Let $n < N$ be coprime to p and $c \in R^*$. Then there exists $v \in R^*$ such that $(1 + \pi^n c, v\pi) = 1$.

Proof. Since n is coprime to p, there exists an integer m such that $nm \equiv 1$ modulo p. We have

$$1 = (1 + \pi^n c, -\pi^n c)^m = (1 + \pi^n c, (-1)^m \pi^{nm} c^m) = (1 + \pi^n c, v\pi)$$

with $v = (-1)^m c^m$. □

Lemma 1.4. Let $n = mp < N$ and $i > 0$. Let $u \in R^*$ be such that $\overline{u} \notin \kappa^P$. Suppose $n + i$ is coprime to p. Then every element in $br(K)_{n+i} \setminus br(K)_{n+i+1}$ can be represented by $(1 + u\pi^n, u')$ for some $u' \in R^*$.

Proof. Fix

$$\alpha \in br(K)_{n+1} \setminus br(K)_{n+i+1}.$$

By (1.1) there is an $x \in R^*$ such that

$$\alpha \equiv (1 + x\pi^{n+i}, u) \pmod{br(K)_{n+i+1}}.$$

Let

$$u' = (-1)^{p+1}(u - x\pi^i)(1 + x\pi^{n+i})^{-1}.$$

Since $i > 0$, $u \equiv u'$ modulo π and hence $u' = vu$ for some $v \in R^*$ with $v \equiv 1$ modulo π. Thus, by ([CT], 4.1.1(b)),

$$(1 + x\pi^{n+i}, u) \equiv (1 + x\pi^{n+i}, u') \pmod{br(K)_{n+i+1}}.$$

Since $(1 - z^p y, y) = 0$, we have

$$(1 + x\pi^{n+i}, u') = ((1 + x\pi^{n+i})(1 - (-1)^p \pi^{mp} u'), u').$$

On the other hand, we know

$$(1 + x\pi^{n+i})(1 - (-1)^p \pi^{mp} u') = 1 + u\pi^n.$$

Thus

$$\alpha \equiv (1 + u\pi^n, u') \pmod{br(K)_{n+i+1}},$$

as desired. □

Lemma 1.5. (Saltman [S4]) Let $\alpha \in br(K)_0 \setminus br(K)_1$. Then there exist $u, v \in R^*$ such that $\overline{u} \notin \kappa^p$ and

$$\alpha - (u, v\pi)$$

is unramified.

Proof. First we show by induction that for each $0 \leq i < N$, there exist $u_i, v_i \in R^*$ such that

$$\alpha - (u_i, v_i\pi) \in br(K)_{i+1}$$

with $\overline{u}_i \notin \kappa^p$. By (1.1),

$$\alpha - (u_0, \pi) \in br(K)_1$$

for some $u_0 \in R^*$ with $\overline{u}_0 \notin \kappa^p$. Suppose there exist $u_i, v_i \in R^*$ such that

$$\alpha - (u_i, v_i\pi) \in br(K)_{i+1}$$

and $\overline{u}_i \notin \kappa^p$.

Suppose that $i+1$ is coprime to p. Since $\overline{u}_i \notin \kappa^p$, by (1.1), there exists $c \in R^*$ such that

$$\alpha - (u_i, v_i\pi) - (u_i, 1 + \pi^{i+1}c) \in br(K)_{i+2}.$$

Thus

$$\alpha - (u_i, v_i\pi(1 + \pi^{i+1}c)) \in br(K)_{i+2}.$$

Suppose that $i+1$ is divisible by p and $i+1 < N$. By (1.1), there exists $c \in R^*$ such that

$$\alpha - (u_i, v_i\pi) - (1 + \pi^{i+1}c, v_i\pi) \in br(K)_{i+2}.$$

Then

$$\alpha - (u_i(1 + \pi^{i+1}c), v_i\pi) \in br(K)_{i+2},$$

and $\overline{u_i(1 + \pi^{i+1}c)} = \overline{u}_i$, so the induction hypothesis is confirmed.

In particular,

$$\alpha - (u_{N-1}, v_{N-1}\pi) \in br(K)_N.$$

By (1.1),

$$\alpha - (u_{N-1}, v_{N-1}\pi) = \alpha' + (u, v_{N-1}\pi)$$

with α' unramified and $u \in 1 + \pi R$. Thus

$$\alpha - (u_{N-1}u, v_{N-1}\pi) = \alpha'$$

is unramified and $\overline{u_{N-1}u} \notin \kappa^p$, as desired. □

Lemma 1.6. (Saltman [S4]) Let $1 \leq n < N$ be coprime to p and $\alpha \in br(K)_n \setminus br(K)_{n+1}$. Then there exist $u, v \in R^*$ such that

$$\alpha - (1 + v\pi^n, u\pi)$$

is unramified.

Proof. First we show by induction that for each $0 \leq i < N - n$, there exist $u_i, v_i \in R^*$ such that

$$\alpha - (1 + v_i\pi^n, u_i) \in br(K)_{n+i+1}$$

with $\overline{u}_i \notin \kappa^p$. Since n is coprime to p, by (1.1),

$$\alpha - (1 + v_0\pi^n, u_0) \in br(K)_{n+1}$$

for some $u_0, v_0 \in R^*$ with $\overline{u}_0 \notin \kappa^p$. Suppose there exist $u_i, v_i \in R^*$ such that

$$\alpha - (1 + v_i\pi^n, u_i) \in br(K)_{n+i+1} \setminus br(K)_{n+i+2}$$

and $\overline{u}_i \notin \kappa^p$. We will break the proof into two cases, depending upon the divisibility properties of $n + i + 1$.

Case 1: $n + i + 1$ is coprime to p. Since $\overline{u}_i \notin \kappa^p$, by (1.1), there exists $c \in R^*$ such that

$$\alpha - (1 + v_i\pi^n, u_i) - (1 + \pi^{n+i+1}c, u_i) \in br(K)_{n+i+2}.$$

Let

$$v_{i+1} = v_i + \pi^{i+1}c + \pi^{n+i+1}v_ic.$$

Then

$$\alpha - (1 + v_{i+1}\pi^n, u_i) \in br(K)_{n+i+2},$$

verifying the inductive hypothesis.

Case 2: $n + i + 1$ is divisible by p and $n + i + 1 < N$. Since n is coprime to p, by (1.3) there is a $v \in R^*$ such that

$$(1 + v_i\pi^n, v\pi) = 1.$$

In particular,

$$(1 + v_i\pi^n, u_i) = (1 + v_i\pi^n, u_iv\pi).$$

By (1.1), there exists $c \in R^*$ such that

$$\alpha - (1 + v_i\pi^n, u_i) - (1 + \pi^{n+i+1}c, u_iv\pi) \in br(K)_{n+i+2}.$$

Thus

$$\begin{aligned}\alpha - (1 + v_i\pi^n, u_i) - (1 + \pi^{n+i+1}c, u_iv\pi) &= \alpha - (1 + v_i\pi^n, u_iv\pi) - (1 + \pi^{n+i+1}c, u_iv\pi) \\ &= \alpha - ((1 + v_i\pi^n)(1 + \pi^{n+i+1}c), u_iv\pi) \\ &\in br(K)_{n+i+2}.\end{aligned}$$

Let $v_{i+1} = v_i + \pi^{i+1}c + v_ic\pi^{n+i+1}$. Then

$$\alpha - (1 + v_{i+1}\pi^n, u_iv\pi) \in br(K)_{n+i+2}.$$

Once again, by (1.3),

$$(1 + v_{i+1}\pi^n, v'\pi) = 1$$

for some $v' \in R^*$. Hence

$$(1 + v_{i+1}\pi^n, v'^{-1}\pi^{-1}) = 1$$

and

$$\begin{aligned}(1 + v_{i+1}\pi^n, u_iv\pi) &= (1 + v_{i+1}\pi^n, u_iv\pi)(1 + v_{i+1}\pi^n, v'^{-1}\pi^{-1}) \\ &= (1 + v_{i+1}\pi^n, u_ivv'^{-1}).\end{aligned}$$

Let $u_{i+1} = u_ivv'^{-1}$. Then

$$\alpha - (1 + v_{i+1}\pi^n, u_{i+1}) \in br(K)_{n+i+2}.$$

Suppose that $\overline{u}_{i+1} \in \kappa^p$. Replacing u_{i+1} by a^pu_{i+1}, we assume that $\overline{u}_{i+1} = 1$. Then, by ([CT], 4.1.1(b)), $(1 + v_{i+1}\pi^n, u_{i+1}) \in br(K)_{n+1}$, contradicting the fact that $\alpha \notin br(K)_{n+1}$. Thus $\overline{u}_{i+1} \notin \kappa^p$, thereby verifying the inductive hypothesis.

In particular, for $i = N - n$, we have

$$\alpha - (1 + v_{N-1}\pi^n, u_{N-1}) \in br(K)_N.$$

Since n is coprime to p, by (1.3), we have

$$(1 + v_{N-1}\pi^n, u_{N-1}) = (1 + v_{N-1}\pi^n, u_{N-1}v\pi)$$

for some $v \in R^*$. By (1.1(d)),

$$\alpha - (1 + v_{N-1}\pi^n, u_{N-1}v\pi) = \alpha' + (1 + x'b^p, u_{N-1}v\pi)$$

for some α' unramified and $x' \in R$. Hence

$$\alpha - ((1 + v_{N-1}\pi^n)(1 + x'b^p), u_{N-1}v\pi) = \alpha'$$

is unramified, as desired. □

Lemma 1.7. (Saltman [S4]) Let $1 < n < N$ be divisible by p and

$$\alpha \in br(K)_n \setminus br(K)_{n+1}.$$

Then there exist $u, v \in R^*$ such that

$$\alpha - (1 + u\pi^n, v\pi)$$

is unramified.

Proof. First we show by induction that for each $0 \leq i < N - n$, there exist $u_i, v_i \in R^*$ such that

$$\alpha - (1 + u_i\pi^n, v_i\pi) \in br(K)_{n+i+1}$$

and

$$\overline{u}_i \notin \kappa^p.$$

Since n is divisible by p, by (1.1) there exists $u_0 \in R^*$ with $\overline{u}_0 \notin \kappa^p$ such that

$$\alpha - (1 + u_0\pi^n, \pi) \in br(K)_{n+1}.$$

Suppose there exist $u_i, v_i \in R^*$ such that

$$\alpha - (1 + u_i\pi^n, v_i\pi) \in br(K)_{n+i+1} \setminus br(K)_{n+i+2}$$

and $\overline{u}_i \notin \kappa^p$. We again break into two cases.

Case 1: $n + i + 1$ is divisible by p. By (1.1), there exists $c \in R^*$ such that

$$\alpha - (1 + u_i\pi^n, v_i\pi) - (1 + \pi^{n+i+1}c, v_i\pi) \in br(K)_{n+i+2}.$$

Thus

$$\begin{aligned}\alpha - ((1 + u_i\pi^n)(1 + \pi^{n+i+1}c), v_i\pi) &= \alpha - (1 + (u_i + \pi^{i+1}c + u_i\pi^{n+i+1}c)\pi^n, v_i\pi) \\ &\in br(K)_{n+i+2}.\end{aligned}$$

Let

$$u_{i+1} = u_i + \pi^{i+1}c + u_i\pi^{n+i+1}c.$$

Then $\overline{u}_{i+1} = \overline{u}_i \notin \kappa^p$ and

$$\alpha - (1 + u_{i+1}\pi^n, v_i\pi) \in br(K)_{n+i+2},$$

establishing the inductive hypothesis.

Case 2: $n+i+1$ is coprime to p. By (1.1), there are $c, f \in R^*$ with $\overline{f} \notin \kappa^p$ such that

$$\alpha - (1+u_i\pi^n, v_i\pi) - (1+c\pi^{n+i+1}, f) \in br(K)_{n+i+2}.$$

By (1.4), there is an $f' \in R^*$ such that

$$(1+c\pi^{n+i+1}, f) = (1+v_i\pi^n, f').$$

In particular

$$\begin{aligned}\alpha - (1+u_i\pi^n, v_if'\pi) &= \alpha - (1+u_i\pi^n, v_i\pi) - (1+u_i\pi^n, f') \\ &= \alpha - (1+u_i\pi^n, v_i\pi) - (1+c\pi^{n+i+1}, f) \\ &\in br(K)_{n+i+2}.\end{aligned}$$

Proceding as in the proof of (1.5), we conclude that there are $u, v \in R^*$ such that

$$\alpha - (1+u\pi^n, v\pi)$$

is unramified. □

Proposition 1.8. (Saltman [S4]) Let $\alpha \in {}_p\mathrm{Br}(K)$. Then there exists a parameter π' such that $\alpha \otimes K(\sqrt[p]{\pi'})$ is unramified.

Proof. If $\alpha \in br(K)_n$ for $0 \leq n < N$ $\alpha \in br(K)_N$, then applying (1.5, 1.6, 1.7) we see that there is a uniformizer π and an element $a \in K$ such that

$$\alpha - (a, \pi)$$

is unramified. If $\alpha \in br(K)_N$ then by (1.1) we have (in fact) that there is an element $x \in R$ such that

$$\alpha - (1+x\pi^N, \pi)$$

is unramified for any choice of uniformizer π. In any case, this implies that for the appropriate choice of π, the base change

$$\alpha \otimes K(\sqrt[p]{\pi})$$

is unramified, as desired. □.

2. The Brauer group of the function field of a surface

Let R be an integral domain with field of fractions K. Let C be a central simple algebra over K.

Definitions.

- The algebra C is *unramified* on R if there is an Azumaya algebra $\mathcal{C}$ over R such that $\mathcal{C} \otimes_R K$ is Brauer equivalent to C.
- An element in $\mathrm{Br}(K)$ is *unramified* on R if it is represented by central simple algebra over K which is unramified on R.
- If an element in $\mathrm{Br}(K)$ is not unramified on R, then we say that it is *ramified* on R.
- Let P be a prime ideal of R. If an element in $\mathrm{Br}(K)$ is unramified (resp. ramified) on R_P, then we say that α is unramified (resp. ramified) at P.

Let X be a regular integral scheme and K its function field. Let X^i be the set of points of X of codimension i. For $x \in X^i$, let $\mathcal{O}_{X,x}$ be the local ring at x, m_x be the maximal ideal at x and $\kappa(x)$ be the residue field at x. For $x \in X^1$, if a central simple algebra C over K is unramified (resp. ramified) at $\mathcal{O}_{X,x}$, then we say that C is unramified (resp. ramified) at x. Let K_x denote the field of fractions of the completion of $\mathcal{O}_{X,x}$ at m_x. If m_x is generated by π, we also denote K_x by K_π. For $x \in X^1$, let ν_x be the valuation on K given by x. Suppose that $\mathrm{char}(K) = 0$ and $\mathrm{char}(\kappa(x)) = p > 0$. Let $N_x = \nu_x(p)p/(p-1)$. If m_x is generated by π, we also denote N_x by N_π.

We will make (implicit) essential use of the following results throughout the rest of this paper. The following results are true in much more generality for torsors (cf. [CTS], 6.13, [APS], 4.3).

Lemma 2.1 Suppose X is a Noetherian two dimensional regular integral scheme with function field K. Given $\alpha \in \mathrm{Br}(K)$, the following are equivalent.

1. For every discrete valuation of K with center on X and valuation ring R, α is unramified on R.

2. For every discrete valuation of K with center of codimension 1 on X and valuation ring R, α is unramified on R.

3. For any central simple algebra C over K representing α, there is a sheaf of Azumaya algebras A on X and an isomorphism $A \otimes_{\mathscr{O}_X} K \xrightarrow{\sim} C$ of K-algebras.

Proof. Fix a central simple algebra C over K representing α and let A be a maximal order in C. By [Prop. 1.2, AG] any localization of A at a point of X is a maximal order in C over $\mathscr{O}_{X,x}$. Since X is regular and 2-dimensional and A is reflexive as an R-module [Thm. 1.5, AG], A is a finite locally free R-algebra. Consider the map

$$\mu : A \otimes A^\circ \to \mathscr{E}nd_{\mathscr{O}_X}(A)$$

of locally free sheaves of $\mathscr{O}_X$-modules of rank n^2 associated to the map of presheaves that sends an elementary tensor $a \otimes b$ in $A(U) \otimes_{\mathscr{O}(U)} A(U)$ to the $\mathscr{O}_X(U)$-linear map sending x to axb. The maximal order A is Azumaya if and only if μ is an isomorphism. We therefore assume that $X = \mathrm{Spec}(B)$ with B a regular local ring of dimension 2. In this case $A \otimes_B A^\circ$ and $End_B(A)$ are free B-modules of same rank, since $\mu \otimes_B K$ is an isomorphism. For a choice of a basis of these modules over B, μ is an isomorphism if and only if the determinant of μ is a unit in B. By Krull's theorem, the determinant is a unit in B if and only if each it is a unit at every codimension one point. This shows that the second two conditions are equivalent.

On the other hand, the first condition certainly implies the second, and hence the third. It remains to show that the third condition implies the first. Suppose A is an Azumaya algebra on X restricting to C. If R is a valuation ring with center on X then there is a commutative diagram



in which the two diagonal arrows are the canonical ones. Pulling A back yields an Azumaya algebra on R restricting to C on K, showing that C is unramified on R, as desired. □

Corollary 2.2 Suppose that R is a two dimensional regular Noetherian integral domain with field of fractions K. Then a central simple algebra C over K is unramified on R if and only if it is unramified on R_P for every height one prime ideal P of R.

Corollary 2.3 Suppose that X is a Noetherian two dimensional regular scheme with fraction field K. Then an element $\alpha \in \mathrm{Br}(K)$ is unramified if and only if it is in the image of the injective restriction map

$$\mathrm{Br}(X) \to \mathrm{Br}(K).$$

Lemma 2.4 Let R be a discrete valuation ring with field of fractions K. Let $\hat{R}$ be the completion of R at the discrete valuation and $\hat{K}$ the field of fractions of $\hat{R}$. Then a central simple algebra C over K is unramified at R if and only if $C \otimes_K \hat{K}$ is unramified at $\hat{R}$.

Proof. Let $\mathcal{A}$ be an Azumaya algebra over $\hat{R}$ such that there is an isomorphism $\phi : \mathcal{A} \otimes_{\hat{R}} \hat{K} \to C \otimes_K \hat{K}$. Then $\mathcal{C} = \mathcal{A} \times_\phi C = \{(a, b) \in \mathcal{A} \times C \mid \phi(a \times 1) = (b, 1)\}$ is an Azumaya R-algebra with $\mathcal{C} \otimes_R K \simeq C$. □

Lemma 2.5 Let R be a discrete valuation ring with field of fractions K and residue field F. Suppose that $\mathrm{char}(F) \neq p$ and $\alpha \in {}_p\mathrm{Br}(K)$. If π is a parameter of R, $\alpha \otimes_K K(\sqrt[p]{\pi})$ is unramified on $R[\sqrt[p]{\pi}]$.

Proof. See ([S2], 0.4). □

Proposition 2.6. Let A be a regular local ring of dimension two with maximal ideal $m = (\pi, \delta)$. Let K be the field of fractions of A and k the residue field of A. Suppose that $\mathrm{char}(K) = 0$, $\mathrm{char}(k) = p > 0$ and $k = k^p$. Let $\alpha \in {}_p\mathrm{Br}(K)$. If α is ramified on A at most at (π) and (δ), then $\alpha \otimes_K K(\sqrt[p]{\pi}, \sqrt[p]{\delta})$ is unramified at any discrete valuation of $K(\sqrt[p]{\pi}, \sqrt[p]{\delta})$ dominating A.

Proof. Let ν be a discrete valuation of $L = K(\sqrt[p]{\pi}, \sqrt[p]{\delta})$ dominating A. Let R be the valuation ring at ν. The completion of A at m is contained in the

completion of R at its maximal ideal. By (2.4), to show that α is unramified at R, we replace A by its completion at m and assume that A is complete.

Let $B = A[\sqrt[p]{\pi}, \sqrt[p]{\delta}]$. Then B is a complete regular local ring with field of fractions L and B is the integral closure of A in L. Hence $B \subset R$. Thus, to show that α is unramified at ν, it is enough to show that α is unramified on B. Since B is a regular ring of dimension 2, by (2.2), it is enough to show that α is unramified at every height one prime ideal of B.

Let Q be a height one prime ideal of B. Since B is integral over A, $Q \cap A = P$ is a height one prime ideal of A. Suppose $P \neq (\pi)$ and $P \neq (\delta)$. Then α is unramified at P and hence α is unramified at Q.

Suppose $P = (\pi)$. Then $L_Q = K_P(\sqrt[p]{\pi}, \sqrt[p]{\delta})$. Suppose that $\mathrm{char}(\kappa(P)) \neq p$. Since π is a parameter at P, by (2.5), α is unramified over $K_P(\sqrt[p]{\pi}) \subset L_Q$ and hence unramified at Q. Suppose that $\mathrm{char}(\kappa(P)) = p$. Since A is a complete regular local ring of dimension 2 with maximal ideal $m = (\pi, \delta)$, we have $A/P \simeq k[[\overline{\delta}]]$, where $\overline{\delta}$ is the image of δ in A/P. Since $k = k^p$, $[\kappa(P) : \kappa(P)^p] = p$ and $\overline{\delta}$ is not a p^{th} power in $\kappa(P)$. Thus, by (1.2), α is split over $K_Q = K_P(\sqrt[p]{\pi}, \sqrt[p]{\delta})$. In particular, α is unramified at Q. The case $P = (\delta)$ is similar. □

Proposition 2.7. Let A be a regular local ring of dimension two with maximal ideal m and $\pi \in m \setminus m^2$. Let K be the field of fractions of A and k the residue field of A. Suppose that $\mathrm{char}(K) = 0$ and $\mathrm{char}(k) = p > 0$. Let $\alpha \in {}_p\mathrm{Br}(K)$. Suppose α is ramified on A at most at (π). Further assume that α is unramified over $K_\pi(\sqrt[p]{\pi})$. Then α is unramified over any discrete valuation of $K(\sqrt[p]{\pi})$ dominating A.

Proof. Let $B = A[\sqrt[p]{\pi}]$. Then B is a regular local ring with field of fractions $K(\sqrt[p]{\pi})$ and B is integral over A. Every discrete valuation of $L = K(\sqrt[p]{\pi})$ dominating A also dominates B. Thus, B being regular, it is enough to show that α is unramified on B. Let Q be a height one prime ideal of B. Since B is integral over A, $Q \cap A = P$ is a height one prime ideal. Suppose $P \neq (\pi)$. Since α is unramified at P, α is unramified at Q. Suppose $P = (\pi)$. Then by the assumption, α is unramified over $L_Q = K_\pi(\sqrt{\pi})$ and hence unramified at Q by (2.4). □

Lemma 2.8. Let A be a one-dimensional Noetherian local domain with field of fractions K and residue field k. Suppose that $\mathrm{char}(K) = p > 0$ and

$[K : K^p] \leq p^{d+1}$ for some $d \geq 0$. Then $[k : k^p] \leq p^d$.

Proof. Let B be the integral closure of A in K and m a maximal ideal of B. Then B/m is a finite extension of k and hence $[(B/m) : (B/m)^p] = [k : k^p]$. Thus, by replacing A by B_m, we assume that A is a discrete valuation ring. Let $\pi \in A$ be a generator of the maximal ideal of A. Let $u_1, \cdots, u_r \in A^*$ be such that the images of $u_1, \cdots, u_r$ in k are p-independent. Then, it is easy to see that $\pi, u_1, \cdots, u_r$ are p-independent in K. □

Theorem 2.9. Let X an excellent regular integral scheme of dimension 2 and K the function field of X. Let p be a prime number. Suppose that $\mathrm{char}(K) = 0$ and K contains a primitive p^{th} root of unity. Suppose that for every codimension one point x of X with $\mathrm{char}(\kappa(x)) = p$, $[\kappa(x) : \kappa(x)^p] = p$. Let $\alpha \in {}_p\mathrm{Br}(K)$. Then there exist $f, g, h \in K^*$ such that $\alpha \otimes K(\sqrt[p]{f}, \sqrt[p]{g}, \sqrt[p]{h})$ is unramified at every discrete valuation of $K(\sqrt[p]{f}, \sqrt[p]{g}, \sqrt[p]{h})$ whose restriction to K is centered at a point of X.

Proof. Let P be a closed point of X. Suppose that $\mathrm{char}(\kappa(P)) = p$. Let $\pi \in \mathcal{O}_{X,P}$ be a prime dividing p. Then $A = \mathcal{O}_{X,P}/(\pi)$ is a one-dimensional Noetherian local domain. Let $\kappa(\pi)$ be the field of fractions of A. Then $\kappa(\pi)$ is the residue field of a codimension one point of X and $\mathrm{char}(\kappa(\pi)) = p$. Thus, by the assumption, $[\kappa(\pi) : \kappa(\pi)^p] = p$. Hence, by (2.8), $[\kappa(P) : \kappa(P)^p] = 1$. Let X' be a blow-up of X at finitely many closed points. Let $x' \in X'$ be a point of codimension 1. If x' is an exceptional curve, then $\kappa(x') = \kappa(P)(t)$ for some closed point P of X and t a variable. In particular, $[\kappa(x') : \kappa(x')^p] = p$. Thus X' also satisfies the hypothesis of the theorem.

By blowing up X at finitely many closed points ([Li]), we assume that $\mathrm{ram}(\alpha)$ and $\mathrm{Supp}(p)$ are contained in $C+E$, where C and E are regular curves with C intersecting E transversally. Let D be an irreducible component in $C \cup E$. If $\mathrm{char}(\kappa(D)) \neq p$, let π_D be any parameter at D. Then, by (2.5), α is unramified over $K_D(\sqrt[p]{\pi_D})$. If $\mathrm{char}(\kappa(D)) = p$, then by the assumption, $[\kappa(D) : \kappa(D)^p] = p$ and hence by (1.8) there exists a parameter $\pi_D \in K_D^*$ at D such that α is unramified over $K_D(\sqrt[p]{\pi_D})$. By the weak approximation, choose $f \in K^*$ such that $f = \pi_D$ modulo K_D^{*p} for every irreducible component D in $C \cup E$. Then, we have $K_D(\sqrt[p]{f}) = K_D(\sqrt[p]{\pi_D})$ and

$$div(f) = C + E + F$$

for some F not containing any component of C and E. In particular, α is unramified over $K_D(\sqrt[p]{f})$ for every D in $C \cup E$.

Let $\mathcal{P}$ be the finite set of closed points containing $C \cap E$, $C \cap F$ and $E \cap F$ and at least one point from each component of C, E, F. Let A be the semi-local regular two dimensional ring at the points in $\mathcal{P}$. Let $\pi, \delta \in A$ be such that the divisor of π on A is C and the divisor of δ on A is E. For $P \in \mathcal{P}$, let m_P denote the maximal ideal of A at P. If $P \in C \cap F$, $P \notin E$, let $\delta_P \in m_P$ be such that $\delta_P \notin m_Q$ for all $Q \in \mathcal{P}, Q \neq P$ and $\delta_P \notin (\pi) + m_P^2$. Similarly, choose π_Q for each $Q \in E \cap F$, $Q \notin C$.

Let

$$g = \pi \prod_{Q \in (E \cap F) \setminus C} \pi_Q \quad \text{and} \quad h = \delta \prod_{P \in (C \cap F) \setminus E} \delta_P.$$

Let $P \in \mathcal{P}$. By the choice of g and h, we have $m_P = (g, h)$. If $P \in C$, then g defines C at P and if $P \in E$, then h defines E at P.

We claim that α is unramified over $L = K(\sqrt[p]{f}, \sqrt[p]{g}, \sqrt[p]{h})$. Let ν be a discrete valuation of L. Let R be the discrete valuation ring of L at ν. Then there is a point x of X such that R dominates the local ring $A_x = \mathcal{O}_{X,x}$.

If x is not on $C \cup E$, then α is unramified on A_x and hence unramified at ν. Suppose that $x \in C \cup E$.

Suppose that x is a codimension one point. Then x corresponds to an irreducible component of C or E. By the choice of f, α is unramified over $K_x(\sqrt[p]{f}) \subset L_\nu$. In particular α is unramified at ν.

Suppose that x is a closed point. Suppose that $x \notin \mathcal{P}$. Since $x \in C \cup E$, C, E are regular curves on X, $f \in m_x \setminus m_x^2$ and α is ramified on $\mathcal{O}_{X,x}$ only along (f). By the choice of f and by (2.7), α is unramified at ν restricted to $K(\sqrt[p]{f})$. Since $K(\sqrt[p]{f}) \subset L$, α is unramified at ν.

Assume that $x \in \mathcal{P}$. Then, by the choice of g and h, the maximal ideal m_x of $\mathcal{O}_{X,x}$ is generated by g and h. Further α is unramified on A except at (g) and (h). Since ν is centered on $\mathcal{O}_{X,x}$, its restriction to $K(\sqrt[p]{g}, \sqrt[p]{h})$ is also centered on $\mathcal{O}_{X,x}$. Hence, by (2.6), α is unramified at ν restricted to $K(\sqrt[p]{g}, \sqrt[p]{h})$. Since $K(\sqrt[p]{g}, \sqrt[p]{h}) \subset L$, α is unramified at ν. □

Corollary 2.10. Let k be a p-adic field and K a function field of a curve over k. Then for every central simple algebra A over K, index(A) divides $(\text{period}(A))^3$.

Proof. Let A be a central simple algebra over K and α its class in $Br(K)$. Let n be the period of A. It is enough to prove the result for a prime n and assuming K contains a primitive n^{th} root of unity. If n is coprime to p, then by ([S1]), period of A divides the square of the index. Assume that $n = p$. Since K is a field of fractions of a regular proper scheme over the ring of integers in k of dimension 2, by (2.9), there exist $f, g, h \in K^*$ such that α is unramified over $K(\sqrt[p]{f}, \sqrt[p]{g}, \sqrt[p]{h})$. Since the unramified Brauer group of $K(\sqrt[p]{f}, \sqrt[p]{g}, \sqrt[p]{h})$ is zero ([G], 2.15 and 3.1), α is trivial over $K(\sqrt[p]{f}, \sqrt[p]{g}, \sqrt[p]{h})$. Hence the index of α divides p^3. □.

Corollary 2.11. Let k be a number field and K function field of a curve over K. Then for every central simple algebra of period p, there exist $f, g, h \in K^*$ such that $\alpha \otimes K(\sqrt[p]{f}, \sqrt[p]{g}, \sqrt[p]{h})$ is unramified at every discrete valuation of $K(\sqrt[p]{f}, \sqrt[p]{g}, \sqrt[p]{h})$.

Proof. We assume that K contains a primitive p^{th} root of unity. Since K is a field of fractions of a regular proper scheme over the ring of integers in k of dimension 2 and every discrete valuation of K is centered at a point of X, by (2.9), there exist $f, g, h \in K^*$ such that α is unramified over $K(\sqrt[p]{f}, \sqrt[p]{g}, \sqrt[p]{h})$ at every discrete valuation of $K(\sqrt[p]{f}, \sqrt[p]{g}, \sqrt[p]{h})$. □

3. Period-index bounds

In this section we ameliorate some of the results of [L2] on the period-index problem for Brauer classes on arithmetic surfaces. Let k be a number field, $\mathbf{A}$ the ring of adèles of k and S the scheme of integers of k. We fix a regular proper surface X with a flat projective generically smooth morphism $X \to S$. We will write C for the generic fiber of X/S, and we assume that C is geometrically connected. Unlike in Section 7 of [L2], we do not assume that C admits a k-point; instead, we will use the following Lemma to get 0-cycles of degree 1 on appropriate moduli spaces.

Lemma 3.1 Let $\overline{M}$ be a proper smooth connected k-scheme. For every nonempty open subscheme $M \subset \overline{M}$, we have that M contains a 0-cycle of degree 1 if and only if $\overline{M}$ contains a 0-cycle of degree 1.

Proof. It is enough to show that if $\overline{M}$ contains a point with residue field a finite extension L/k then M contains a 0-cycle of degree equal to $[L:k]$. Since $\overline{M}$ is smooth, it is easy to see (by choosing generic parameters in the local ring at Q and taking a Zariski closure) that there is a smooth projective curve D over k with an L-rational point $R \in D$ and a finite morphism $f : D \to \overline{M}$ such that $f(R) = Q$ and $U = f^{-1}(M) \neq \emptyset$. Since any point of U is ample, it is easy to see that there is a divisor $E \subset U \subset D$ of degree $[L:k]$. The proper pushforward of E to $\overline{M}$ gives a 0-cycle of degree equal to $[L:k]$ contained in M, as desired. □

Fix a prime number ℓ and a μ_ℓ-gerbe $\mathscr{X} \to X$. Write $\mathscr{C} \to C$ for the restriction of $\mathscr{X}$ to C. Let $\mathscr{M} := \mathscr{M}_{\mathscr{C}}(\ell)$ denote the stack of stable $\mathscr{C}$-twisted sheaves of rank ℓ and trivial determinant (as described in [L1]). We know the following facts about this stack.

1. $\mathscr{M}$ is a μ_ℓ-gerbe over a smooth quasi-projective variety M admitting a locally factorial compactification $M \subset M^{ss}$ such that

$$\operatorname{codim}(M^{ss} \setminus M, M^{ss}) > 2$$

and

$$\operatorname{Pic}(M^{ss}) = \operatorname{Pic}(M) = \mathbf{Z}.$$

2. the Brauer group $\operatorname{Br}(M \otimes \overline{k})$ is generated by the class associated to $\mathscr{M} \otimes \overline{k}$, this class has period and index ℓ, and the sequence

$$0 \to \operatorname{Br}(k) \to \operatorname{Br}(M) \to \operatorname{Br}(M \otimes \overline{k}) \to 0$$

is exact. Note that since $\mathscr{M}$ is a μ_ℓ-gerbe over M, its Brauer class has period ℓ and thus the universal obstruction provides a *canonical* splitting of the above exact sequence.

Proposition 3.2 The Brauer-Manin set $M(\mathbf{A})^{\mathrm{Br}}$ is non-empty.

Proof. We first claim that for every place v of k, the category $\mathscr{M}(k_v)$ is non-empty. Indeed, consider the base change $\mathscr{X} \otimes_S \mathscr{O}_{k_v} \to X \otimes_S \mathscr{O}_{k_v}$. This is a μ_ℓ-gerbe on a proper curve over a complete discrete valuation ring with finite residue field. Since the Brauer group of such a scheme is trivial, we

have an invertible $\mathscr{C}_{k_v}$-twisted sheaf $\mathscr{L}$. Writing $L := (\mathscr{L}^{\otimes \ell})^\vee$, tensoring with $\mathscr{L}$ defines an isomorphism

$$\mathscr{M}_{C_{k_v}}(\ell, L) \xrightarrow{\sim} \mathscr{M} \otimes k_v.$$

Since stable vector bundles of rank ℓ and determinant L exist on any curve over an infinite field, the former has an object over k_v, whence the latter does as well.

Write (x_v) for the system of objects of $\mathscr{M}(k_v)$. Projecting to M gives a point $(\overline{x}_v) \in M(\mathbf{A})$ with the property that the pairing $(\overline{x}_v) \cdot [\mathscr{M} \to M]$ is 0 in $\mathbf{Q}/\mathbf{Z}$. As noted above, the Leray spectral sequence shows that $\mathrm{Br}(M)/\mathrm{Br}(k)$ is generated by the class of $\mathscr{M} \to M$. We conclude that the point $(\overline{x}_v)$ lies in $M(\mathbf{A})^{\mathrm{Br}}$, as desired. □

Theorem 3.3 Assuming the CT-conjecture (see page 2), any class $\alpha \in \mathrm{Br}(X)$ satisfies $\mathrm{ind}(\alpha) | \mathrm{per}(\alpha)^2$.

Proof. By standard inductive arguments, we may assume that α has prime index ℓ.

We retain the notation from above. Let $M \subset \overline{M}$ be any smooth compactification of M; this is possible by Hironaka's theorem, since k has characteristic 0. By functoriality we have that $M(\mathbf{A})^{\mathrm{Br}} \subset \overline{M}(\mathbf{A})^{\mathrm{Br}}$, hence by Propostion 3.2 $\overline{M}(\mathbf{A})^{\mathrm{Br}} \neq \emptyset$. The CT-conjecture applies to show that $\overline{M}$ has a 0-cycle of degree 1. Applying Lemma 3.1, we have a 0-cycle of degree 1 in M. Put another way, there are two finite étale k-algebras A_1 and A_2 such that $[A_1 : k]$ is relatively prime to $[A_2 : k]$ and such that $M(A_1) \neq \emptyset$ and $M(A_2) \neq \emptyset$.

Given a finite étale closed subscheme $Z \in M$, the Brauer class of $\mathscr{M} \times_M Z$ lies in $\mathrm{Br}(Z)[\ell]$ and thus has index dividing ℓ (as the residue fields of the Artinian scheme Z are global fields). It follows that there is a finite flat Z-scheme $Y \to Z$ of degree ℓ such that $\mathscr{M}(Y) \neq \emptyset$. Applying this to A_1 and A_2 above, we find that for $i = 1, 2$ there is a finite flat A_i-algebra B_i of degree ℓ and a locally free $\mathscr{C}_{B_i}$-twisted sheaf of rank ℓ. Putting these together and arguing as in Section 4.1.1 of [L3], we see that the index of α divides $\ell^2[A_i : k]$ for $i = 1, 2$. Since $[A_1 : k]$ and $[A_2 : k]$ are relatively prime, we see that $\mathrm{ind}(\alpha) | \ell^2$, as desired. □

4. u-invariant of the function field a curve over a number field

Theorem 4.1. Assume that the CT-conjecture holds. Let k be a number field and K the function field of a curve over k. Then for any central simple algebra A over K, $\mathrm{ind}(A)$ divides $\mathrm{period}(A)^5$.

Proof. Let A be a central simple algebra over K. By a standard inductive argument, it is enough to consider the case $\mathrm{period}(A) = p$ a prime and K contains a primitive p^{th} root of unity. By (2.11), there exist $f, g, h \in K^*$ such that $A \otimes_K K(\sqrt[p]{f}, \sqrt[p]{g}, \sqrt[p]{h})$ is unramified at every discrete valuation of $K(\sqrt[p]{f}, \sqrt[p]{g}, \sqrt[p]{h})$. By (3.3), $\mathrm{index}(A \otimes K(\sqrt[p]{f}, \sqrt[p]{g}, \sqrt[p]{h}))$ divides p^2. Hence $\mathrm{index}(A)$ divides p^5. □

Theorem 4.2. Assume that the CT-conjecture holds. Let k be a number field and K the function field of a curve over k. Then there exists an integer N_2 (which does not depend on K or k) such that every element in $H^2(K, \mu_2)$ is a sum of at most N_2 symbols.

Proof. Let $\alpha \in H^2(K, \mu_2)$. By (4.1), $\mathrm{index}(\alpha)$ divides 2^5. Let $\mathcal{A}$ be the generic division algebra of degree 2^5 with center Z and $X_{\mathcal{A}^2}$ be the Severi-Brauer variety of $\mathcal{A} \otimes \mathcal{A}$ over Z. Let α be represented by a central simple algebra A over K of degree 2^5. Then there is a specialisation from the function field $Z(X_{\mathcal{A}^2})$ to K which specialises $\mathcal{A}$ to A. Since $\mathcal{A}$ is 2-torsion element in the Brauer group of $Z(X_{\mathcal{A}^2})$, $\mathcal{A}$ is a product of N_2 quaternion algebras over $Z(X_{\mathcal{A}^2})$ ([M]). In particular A is a product of N_2 quaternion algebras over K. Hence α is a sum of at most N_2 symbols. □.

Theorem 4.3. Assume that the CT-conjecture holds. Let k be a number field and K the function field of a curve over k. Then every element in $H^3(K, \mu_2)$ is a sum of N_2 symbols.

Proof. Let $\beta \in H^3(K, \mu_2)$. Then by ([Su]), there exists $f \in K^*$ such that α is zero over $K(\sqrt{f})$. Hence $\beta = (f) \cdot \alpha$ for some $\alpha \in H^2(K, \mu_2)$ ([A], 4.6). By (4.2), $\alpha = (a_1) \cdot (b_1) + \cdots + (a_{N_2}) \cdot (b_{N_2})$. Thus $\beta = (f) \cdot \alpha = (f) \cdot (a_1) \cdot (b_1) + \cdots + (f) \cdot (a_{N_2}) \cdot (b_{N_2})$. □

Theorem 4.4 Assume that the CT-conjecture holds. Let k be a totally imaginary number field and K the function field of a curve over k. Then $u(K)$ is finite.

Proof. By (4.2), there exists N_2 such that every element in $H^2(K, \mu_2)$ is a sum of at most N_2 symbols. By (4.3), every element in $H^3(K, \mu_2)$ is a sum of at most N_2 symbols. Since k is a totally imaginary number field, $H^4(K, \mu_2) = 0$. Hence $u(K)$ is finite (cf. [PS]). □

References

[A] J. K. Arason, *Cohomologische Invarianten quadratischer Formen*, J.Algebra **36** (1975), 448–491.

[AG] M. Auslander, O. Goldman, *Maximal orders*, Trans. Amer. Math. Soc. **97** (1960), 367–409.

[APS] Asher Auel, R. Parimala and V. Suresh, *Quadric surface bundles over surfaces*, arXiv:1207.4105v1

[B] E. Brussel, *On Saltman's p-adic curves papers*, Quadratic forms, linear algebraic groups, and cohomology, 13-39, Dev. Math., **18**, Springer, New York, 2010.

[CT1] J.-L. Colliot-Thélène *Conjectures de type local-global sur image des groupes de Chow dans la cohomologie étale*, Algebraic K -theory (Seattle, WA, 1997), 1-12, Proc. Sympos. Pure Math., **67**, Amer. Math. Soc., Providence, RI, 1999.

[CT2] J.-L. Colliot-Thélène *Cohomologie des corps valués hensliens, d'après K. Kato et S. Bloch*, Algebraic K-theory and its applications (Trieste, 1997), ed. H. Bass, A. O. Kuku, C. Pedrini, World Scientific Publishing, 1999, p. 120-163.

[CTS] J.-L. Colliot-Thélène et J.-J. Sansuc, *Fibrés quadratiques et composantes connexes réelles*, Math. Ann. **244** (1979), 105-134.

[G] A. Grothendieck, *Le groupe de Brauer III*, Dix exposs sur la cohomologie des schmas, Amsterdam, North-Holland, Amsterdam (1968), 88-188.

[K] K. Kato, *Galois cohomology of complete discrete valuation fields*, in Algebraic K-Theory, Springer L. N. M. **967** (1982), 215-238.

[L1] M. Lieblich, *Moduli of twisted sheaves*, Duke Math. J. **138** (2007), 23-118.

[L2] M. Lieblich, *Period and index in the Brauer group of an arithmetic surface*, with an appendix by Daniel Krashen, J. Reine Angew. Math **659** (2011), 1-41.

[L3] M. Lieblich, *Twisted sheaves and the period-index problem*, Compos. Math. **144** (2008), no. 1, 1-31.

[Li] J. Lipman, *Desingularization of two-dimensional schemes*, Ann. Math. **107** (1978), 151207

[M] A.S. Merkurjev, *On the norm residue symbol of degree 2*, Dokl. Akad. Nauk SSSR **261** (1981), 542-547.

[MS] A.S. Merkurjev and A. A. Suslin, *K-cohomology of Severi–Brauer varieties and the norm- residue homomorphism*, Izv. Akad. Nauk SSSR, Ser. Mat., **46**, No. 5, (1982) 1011-1061.

[S1] D.J. Saltman, *Division Algebras over p-adic curves*, J. Ramanujan Math. Soc. **12** (1997), 25-47.

[S2] D.J. Saltman, *Cyclic Algebras over p-adic curves*, J. Algebra **314** (2007) 817-843.

[S3] D.J. Saltman, *Division algebras over surfaces*, J. Algebra **320** (2008), 1543-1585.

[S4] D.J. Saltman, *Bad characteristic*, private notes.

[PS] R. Parimala and V. Suresh, *On the length of a quadratic form*, Algebra and number theory, 147-157, Hindustan Book Agency, Delhi, 2005.

[Su] V. Suresh, *Galois cohomology in degree 3 of function fields of curves over number fields*, J. Number Theory **107** (2004), no. 1, 80-94.

Department of Mathematics, University of Washington, Box 354350 Seattle, WA 98195

Department of Mathematics & Computer Science, Emory University, 400 Dowman Drive NE W401, Atlanta, GA 30322

E-mail addresses: lieblich@math.washington.edu, parimala@mathcs.emory.edu, suresh@mathcs.emory.edu